

ON HERMITE HADAMARD-TYPE INEQUALITIES FOR STRONGLY φ -CONVEX FUNCTIONS

MEHMET ZEKI SARIKAYA

ABSTRACT. In this paper, we introduce the notion of strongly φ -convex functions with respect to $c > 0$ and present some properties and representation of such functions. We obtain a characterization of inner product spaces involving the notion of strongly φ -convex functions. Finally, a version of Hermite Hadamard-type inequalities for strongly φ -convex functions are established.

1. INTRODUCTION

The inequalities discovered by C. Hermite and J. Hadamard for convex functions are very important in the literature (see, e.g., [4], [6, p.137]). These inequalities state that if $f : I \rightarrow \mathbb{R}$ is a convex function on the interval I of real numbers and $a, b \in I$ with $a < b$, then

$$(1.1) \quad f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2}.$$

The inequality (1.1) has evoked the interest of many mathematicians. Especially in the last three decades numerous generalizations, variants and extensions of this inequality have been obtained, to mention a few, see ([3]-[8]) and the references cited therein.

Let us consider a function $\varphi : [a, b] \rightarrow [a, b]$ where $[a, b] \subset \mathbb{R}$. Youness have defined the φ -convex functions in [13]:

Definition 1. A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be φ -convex on $[a, b]$ if for every two points $x \in [a, b], y \in [a, b]$ and $t \in [0, 1]$ the following inequality holds:

$$f(t\varphi(x) + (1-t)\varphi(y)) \leq tf(\varphi(x)) + (1-t)f(\varphi(y)).$$

In [2], Cristescu proved the followig results for the φ -convex functions

Lemma 1. For $f : [a, b] \rightarrow \mathbb{R}$, the following statements are equivalent:

- (i) f is φ -convex functions on $[a, b]$,
- (ii) for every $x, y \in [a, b]$, the mapping $g : [0, 1] \rightarrow \mathbb{R}$, $g(t) = f(t\varphi(x) + (1-t)\varphi(y))$ is classically convex on $[0, 1]$.

Obviously, if function φ is the identity, then the classical convexity is obtained from the previous definition. Many properties of the φ -convex functions can be found, for instance, in [1], [2], [13].

2000 *Mathematics Subject Classification.* 26D10, 26A51, 46C15.

Key words and phrases. Hermite-Hadamard's inequalities, φ -convex functions, strongly convex with modulus $c > 0$.

Recall also that a function $f : I \rightarrow \mathbb{R}$ is called strongly convex with modulus $c > 0$, if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) - ct(1-t)(x-y)^2$$

for all $x, y \in I$ and $t \in (0, 1)$. Strongly convex functions have been introduced by Polyak in [9] and they play an important role in optimization theory and mathematical economics. Various properties and applicatins of them can be found in the literature see ([9]-[12]) and the references cited therein.

In this paper, we introduce the notion of strongly φ -convex functions defined in normed spaces and present some properties of them. In particular, we obtain a representation of strongly φ -convex functions in inner product spaces and, using the methods of [11] and [12], we give a characterization of inner product spaces, among normed spaces, that involves the notion of strongly φ -convex function. Finally, a version of Hermite–Hadamard-type inequalities for strongly φ -convex functions is presented. This result generalizes the Hermite–Hadamard-type inequalities obtained in [10] for strongly convex functions, and for $c = 0$, coincides with the classical Hermite–Hadamard inequalities, as well as the corresponding Hermite–Hadamard-type inequalities for φ -convex functions in [2].

2. MAIN RESULTS

In what follows $(X, \|\cdot\|)$ denotes a real normed space, D stands for a convex subset of X , $\varphi : D \rightarrow D$ is a given function and c is a positive constant. We say that a function $f : D \rightarrow \mathbb{R}$ is strongly φ -convex with modulus c if

$$(2.1) \quad f(t\varphi(x) + (1-t)\varphi(y)) \leq tf(\varphi(x)) + (1-t)f(\varphi(y)) - ct(1-t)\|\varphi(x) - \varphi(y)\|^2$$

for all $x, y \in D$ and $t \in [0, 1]$. We say that f is strongly φ -midconvex with modulus c if (2.1) is assumed only for $t = \frac{1}{2}$, that is

$$f\left(\frac{\varphi(x) + \varphi(y)}{2}\right) \leq \frac{f(\varphi(x)) + f(\varphi(y))}{2} - \frac{c}{4}\|\varphi(x) - \varphi(y)\|^2, \text{ for } x, y \in D.$$

The notion of φ -convex function corresponds to the case $c = 0$. We start with the following lemma which give some relationships between strongly φ -convex functions and φ -convex functions in the case where X is a real inner product space (that is, the norm $\|\cdot\|$ is induced by an inner product: $\|\cdot\| := \langle x|x \rangle$).

Lemma 2. *Let $(X, \|\cdot\|)$ be a real inner product space, D be a convex subset of X and c be a positive constant and $\varphi : D \rightarrow D$.*

i) A function $f : D \rightarrow \mathbb{R}$ is strongly φ -convex with modulus c if and only if the function $g = f - c\|\cdot\|^2$ is φ -convex.

ii) A function $f : D \rightarrow \mathbb{R}$ is strongly φ -midconvex with modulus c if and only if the function $g = f - c\|\cdot\|^2$ is φ -midconvex.

Proof. i) Assume that f is strongly φ -convex with modulus c . Using properties of the inner product, we obtain

$$\begin{aligned}
& g(t\varphi(x) + (1-t)\varphi(y)) \\
&= f(t\varphi(x) + (1-t)\varphi(y)) - c \|t\varphi(x) + (1-t)\varphi(y)\|^2 \\
&\leq tf(\varphi(x)) + (1-t)f(\varphi(y)) - ct(1-t) \|\varphi(x) - \varphi(y)\|^2 - c \|t\varphi(x) + (1-t)\varphi(y)\|^2 \\
&\leq tf(\varphi(x)) + (1-t)f(\varphi(y)) - c \left(t(1-t) \left[\|\varphi(x)\|^2 - 2 \langle \varphi(x) | \varphi(y) \rangle + \|\varphi(y)\|^2 \right] \right. \\
&\quad \left. - \left[t^2 \|\varphi(x)\|^2 + 2t(1-t) \langle \varphi(x) | \varphi(y) \rangle + (1-t) \|\varphi(y)\|^2 \right] \right) \\
&= tf(\varphi(x)) + (1-t)f(\varphi(y)) - ct \|\varphi(x)\|^2 - c(1-t) \|\varphi(y)\|^2 \\
&= tg(\varphi(x)) + (1-t)g(\varphi(y))
\end{aligned}$$

which gives that g is φ -convex function.

Conversely, if g is φ -convex function, then we get

$$\begin{aligned}
f(t\varphi(x) + (1-t)\varphi(y)) &= g(t\varphi(x) + (1-t)\varphi(y)) + c \|t\varphi(x) + (1-t)\varphi(y)\|^2 \\
&\leq tg(\varphi(x)) + (1-t)g(\varphi(y)) + c \|t\varphi(x) + (1-t)\varphi(y)\|^2 \\
&= t \left[g(\varphi(x)) + c \|\varphi(x)\|^2 \right] + (1-t) \left[g(\varphi(y)) + c \|\varphi(y)\|^2 \right] \\
&\quad - ct(1-t) \left[\|\varphi(x)\|^2 - 2 \langle \varphi(x) | \varphi(y) \rangle + \|\varphi(y)\|^2 \right] \\
&= tf(\varphi(x)) + (1-t)f(\varphi(y)) - ct(1-t) \|\varphi(x) - \varphi(y)\|^2
\end{aligned}$$

which shows that f is strongly φ -convex with modulus c .

ii) Assume now that f is strongly φ -midconvex with modulus c . Using the parallelogram law, we have

$$\begin{aligned}
g\left(\frac{\varphi(x) + \varphi(y)}{2}\right) &= f\left(\frac{\varphi(x) + \varphi(y)}{2}\right) - c \left\| \frac{\varphi(x) + \varphi(y)}{2} \right\|^2 \\
&\leq \frac{f(\varphi(x)) + f(\varphi(y))}{2} - \frac{c}{4} \|\varphi(x) - \varphi(y)\|^2 - \frac{c}{4} \|\varphi(x) + \varphi(y)\|^2 \\
&= \frac{f(\varphi(x)) + f(\varphi(y))}{2} - \frac{c}{4} \left(2 \|\varphi(x)\|^2 + 2 \|\varphi(y)\|^2 \right) \\
&= \frac{g(\varphi(x)) + g(\varphi(y))}{2}
\end{aligned}$$

which gives that g is φ -midconvex function.

Similarly, if g is φ -midconvex function, then we get

$$\begin{aligned}
 f\left(\frac{\varphi(x) + \varphi(y)}{2}\right) &= g\left(\frac{\varphi(x) + \varphi(y)}{2}\right) + c \left\| \frac{\varphi(x) + \varphi(y)}{2} \right\|^2 \\
 &\leq \frac{g(\varphi(x)) + g(\varphi(y))}{2} + \frac{c}{4} \|\varphi(x) + \varphi(y)\|^2 \\
 &= \frac{g(\varphi(x)) + \|\varphi(x)\|^2}{2} + \frac{g(\varphi(y)) + \|\varphi(y)\|^2}{2} \\
 &\quad + \frac{c}{4} (\|\varphi(x) + \varphi(y)\|^2 - 2\|\varphi(x)\|^2 - 2\|\varphi(y)\|^2) \\
 &= \frac{f(\varphi(x)) + f(\varphi(y))}{2} - \frac{c}{4} \|\varphi(x) - \varphi(y)\|^2.
 \end{aligned}$$

This completes to proof. $\square$

The following example shows that the assumption that X is an inner product space is essential in the above lemma.

Example. Let $X = \mathbb{R}^2$. Let us consider a function $\varphi : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, defined by $\varphi(x) = x$ for every $x \in \mathbb{R}^2$ and $\|x\| = \max\{|x_1|, |x_2|\}$ for $x = (x_1, x_2)$. Take $f = \|\cdot\|^2$. Then $g = f - \|\cdot\|^2$ is φ -convex being the zero function. However, f is neither strongly φ -convex with modulus 1 nor strongly φ -midconvex with modulus 1. Indeed, for $x = (1, 0)$ and $y = (0, 1)$, we have

$$f\left(\frac{x+y}{2}\right) = \frac{1}{2} \geq \frac{3}{4} = \frac{f(x) + f(y)}{2} - \frac{1}{4} \|x - y\|^2$$

which this contradicts (2.1).

The assumption that X is an inner product space in Lemma 2 is essential. Moreover, it appears that the fact that for every φ -convex function $g : X \rightarrow \mathbb{R}$ the function $f = g + c\|\cdot\|^2$ is strongly φ -convex characterizes inner product spaces among normed spaces. Similar characterizations of inner product spaces by strongly convex and strongly h -convex functions are presented in [11] and [12].

Theorem 1. *Let $(X, \|\cdot\|)$ be a real normed space, D be a convex subset of X and $\varphi : D \rightarrow D$. Then the following conditions are equivalent:*

- i) $(X, \|\cdot\|)$ is a real inner product space;
- ii) For every $c > 0$, $f : D \rightarrow \mathbb{R}$ defined on a convex subset D of X , the function $f = g + c\|\cdot\|^2$ is strongly φ -convex with modulus c ;
- iii) $\|\cdot\|^2 : X \rightarrow \mathbb{R}$ is strongly φ -convex with modulus 1.

Proof. The implication i) $\Rightarrow$ ii) follows by Lemma 2. To see that ii) $\Rightarrow$ iii) take $g = 0$. Clearly, g is φ -convex function, whence $f = c\|\cdot\|^2$ is strongly φ -convex with modulus c . Consequently, $\|\cdot\|^2$ is strongly φ -convex with modulus 1. Finally, to prove iii) $\Rightarrow$ i) observe that by the strongly φ -convexity of $\|\cdot\|^2$, we obtain

$$\left\| \frac{\varphi(x) + \varphi(y)}{2} \right\|^2 \leq \frac{\|\varphi(x)\|^2}{2} + \frac{\|\varphi(y)\|^2}{2} - \frac{1}{4} \|\varphi(x) - \varphi(y)\|^2$$

and hence

$$(2.2) \quad \|\varphi(x) + \varphi(y)\|^2 + \|\varphi(x) - \varphi(y)\|^2 \leq 2\|\varphi(x)\|^2 + 2\|\varphi(y)\|^2$$

for all $x, y \in X$. Now, putting $u = \varphi(x) + \varphi(y)$ and $v = \varphi(x) - \varphi(y)$ in (2.2), we have

$$(2.3) \quad 2 \|u\|^2 + 2 \|v\|^2 \leq \|u + v\|^2 + \|u - v\|^2$$

for all $u, v \in X$.

Conditions (2.2) and (2.3) mean that the norm $\|\cdot\|^2$ satisfies the parallelogram law, which implies, by the classical Jordan-Von Neumann theorem, that $(X, \|\cdot\|)$ is an inner product space. This completes to proof. $\square$

Now, we give a new Hermite–Hadamard-type inequalities for strongly φ -convex functions with modulus c as follows:

Theorem 2. *If $f : [a, b] \rightarrow \mathbb{R}$ is strongly φ -convex with modulus $c > 0$ for the continuous function $\varphi : [a, b] \rightarrow [a, b]$, then*

$$(2.4) \quad \begin{aligned} & f\left(\frac{\varphi(a) + \varphi(b)}{2}\right) + \frac{c}{12} (\varphi(a) - \varphi(b))^2 \\ & \leq \frac{1}{\varphi(b) - \varphi(a)} \int_{\varphi(a)}^{\varphi(b)} f(x) dx \\ & \leq \frac{f(\varphi(a)) + f(\varphi(b))}{2} - \frac{c}{6} (\varphi(a) - \varphi(b))^2. \end{aligned}$$

Proof. From the strongly φ -convexity of f , we have

$$\begin{aligned} f\left(\frac{\varphi(a) + \varphi(b)}{2}\right) &= f\left(\frac{t\varphi(a) + (1-t)\varphi(b)}{2} + \frac{(1-t)\varphi(a) + t\varphi(b)}{2}\right) \\ &\leq \frac{1}{2} f(t\varphi(a) + (1-t)\varphi(b)) + \frac{1}{2} f((1-t)\varphi(a) + t\varphi(b)) - \frac{c}{4} (1-2t)^2 (\varphi(a) - \varphi(b))^2. \end{aligned}$$

By Lemma 1, it follows that the previous inequality can be integrated with respect to t over $[0, 1]$, getting

$$\begin{aligned} & f\left(\frac{\varphi(a) + \varphi(b)}{2}\right) + \frac{c}{12} (\varphi(a) - \varphi(b))^2 \\ & \leq \frac{1}{2} \int_0^1 f(t\varphi(a) + (1-t)\varphi(b)) dt + \frac{1}{2} \int_0^1 f((1-t)\varphi(a) + t\varphi(b)) dt. \end{aligned}$$

In the first integral, we substitute $x = t\varphi(a) + (1-t)\varphi(b)$. Meanwhile, in the second integral we also use the substitution $x = (1-t)\varphi(a) + t\varphi(b)$, we obtain

$$\begin{aligned} & f\left(\frac{\varphi(a) + \varphi(b)}{2}\right) + \frac{c}{12} (\varphi(a) - \varphi(b))^2 \\ & \leq \frac{1}{2(\varphi(a) - \varphi(b))} \int_{\varphi(b)}^{\varphi(a)} f(x) dx + \frac{1}{2(\varphi(b) - \varphi(a))} \int_{\varphi(a)}^{\varphi(b)} f(x) dx \\ & = \frac{1}{\varphi(b) - \varphi(a)} \int_{\varphi(a)}^{\varphi(b)} f(x) dx. \end{aligned}$$

In order to prove the second inequality, we start from the strongly φ -convexity of f meaning that for every $t \in [0, 1]$ one has

$$f(t\varphi(a) + (1-t)\varphi(b)) \leq tf(\varphi(a)) + (1-t)f(\varphi(b)) - ct(1-t)(\varphi(a) - \varphi(b))^2.$$

Lemma 1, allows us to integrate both sides of this inequality on $[0, 1]$, we get

$$\int_0^1 f(t\varphi(a) + (1-t)\varphi(b)) dt \leq f(\varphi(a)) \int_0^1 t dt + f(\varphi(b)) \int_0^1 (1-t) dt - c(\varphi(a) - \varphi(b))^2 \int_0^1 t(1-t) dt.$$

The previous substitution in the first side of this inequality leads to

$$\frac{1}{(\varphi(a) - \varphi(b))} \int_{\varphi(b)}^{\varphi(a)} f(x) dx \leq \frac{f(\varphi(a))}{2} + \frac{f(\varphi(b))}{2} - \frac{c}{6} (\varphi(a) - \varphi(b))^2$$

which gives the second inequality of (2.4). This completes to proof. $\square$

Theorem 3. *If $f : [a, b] \rightarrow \mathbb{R}$ is strongly φ -convex with modulus $c > 0$ for the continuous function $\varphi : [a, b] \rightarrow [a, b]$, then*

$$\begin{aligned} (2.5) \quad & \frac{1}{\varphi(b) - \varphi(a)} \int_{\varphi(a)}^{\varphi(b)} f(x) f(a + b - x) dx \\ & \leq \frac{[f^2(\varphi(x)) + f^2(\varphi(y))]}{6} + \frac{2f(\varphi(x))f(\varphi(y))}{3} \\ & \quad - \frac{c}{6} (\varphi(x) - \varphi(y))^2 [f(\varphi(x)) + f(\varphi(y))] - \frac{c^2}{30} (\varphi(x) - \varphi(y))^4. \end{aligned}$$

Proof. Since f is strongly φ -convex with respect to $c > 0$, we have that for all $t \in [0, 1]$

$$(2.6) \quad f(t\varphi(a) + (1-t)\varphi(b)) \leq tf(\varphi(a)) + (1-t)f(\varphi(b)) - ct(1-t)(\varphi(a) - \varphi(b))^2$$

and

$$(2.7) \quad f((1-t)\varphi(a) + t\varphi(b)) \leq (1-t)f(\varphi(a)) + tf(\varphi(b)) - ct(1-t)(\varphi(a) - \varphi(b))^2.$$

Multiplying both sides of (2.6) by (2.7), it follows that

$$\begin{aligned} (2.8) \quad & f(t\varphi(a) + (1-t)\varphi(b)) f((1-t)\varphi(a) + t\varphi(b)) \\ & \leq t(1-t) [f^2(\varphi(a)) + f^2(\varphi(b))] + (t^2 + (1-t)^2) f(\varphi(a)) f(\varphi(b)) \\ & \quad - ct(1-t)(\varphi(a) - \varphi(b))^2 [f(\varphi(a)) + f(\varphi(b))] + c^2 t^2 (1-t)^2 (\varphi(a) - \varphi(b))^4. \end{aligned}$$

Integrating the inequality (2.8) with respect to t over $[0, 1]$, we obtain

$$\begin{aligned}
& \int_0^1 f(t\varphi(a) + (1-t)\varphi(b))f((1-t)\varphi(a) + t\varphi(b))dt \\
& \leq [f^2(\varphi(a)) + f^2(\varphi(b))] \int_0^1 t(1-t)dt + f(\varphi(a))f(\varphi(b)) \int_0^1 (t^2 + (1-t)^2) dt \\
& \quad - c(\varphi(a) - \varphi(b))^2 [f(\varphi(a)) + f(\varphi(b))] \int_0^1 t(1-t)dt - c^2 (\varphi(a) - \varphi(b))^4 \int_0^1 t^2(1-t)^2 dt \\
& = \frac{[f^2(\varphi(a)) + f^2(\varphi(b))]}{6} + \frac{2f(\varphi(a))f(\varphi(b))}{3} \\
& \quad - \frac{c}{6} (\varphi(a) - \varphi(b))^2 [f(\varphi(a)) + f(\varphi(b))] - \frac{c^2}{30} (\varphi(a) - \varphi(b))^4.
\end{aligned}$$

If we change the variable $x := t\varphi(a) + (1-t)\varphi(b)$, $t \in [0, 1]$, we get the required inequality in (2.5). This proves the theorem. $\square$

Theorem 4. *If $f, g : [a, b] \rightarrow \mathbb{R}$ is strongly φ -convex with modulus $c > 0$ for the continuous function $\varphi : [a, b] \rightarrow [a, b]$, then*

$$\begin{aligned}
(2.9) \quad & \frac{1}{\varphi(b) - \varphi(a)} \int_{\varphi(a)}^{\varphi(b)} f(x)dx \\
& \leq \frac{M(a, b)}{3} + \frac{N(a, b)}{6} - \frac{c}{12} (\varphi(a) - \varphi(b))^2 S(a, b) + \frac{c^2}{30} (\varphi(a) - \varphi(b))^4
\end{aligned}$$

where

$$M(a, b) = f(\varphi(a))g(\varphi(a)) + f(\varphi(b))g(\varphi(b))$$

$$N(a, b) = f(\varphi(a))g(\varphi(b)) + f(\varphi(b))g(\varphi(a))$$

$$S(a, b) = f(\varphi(a)) + f(\varphi(b)) + g(\varphi(a)) + g(\varphi(b)).$$

Proof. Since $f, g : [a, b] \rightarrow \mathbb{R}$ is strongly φ -convex with modulus $c > 0$, we have

$$\begin{aligned}
(2.10) \quad & f(t\varphi(a) + (1-t)\varphi(b)) \leq tf(\varphi(a)) + (1-t)f(\varphi(b)) - ct(1-t)(\varphi(a) - \varphi(b))^2
\end{aligned}$$

$$\begin{aligned}
(2.11) \quad & g(t\varphi(a) + (1-t)\varphi(b)) \leq tg(\varphi(a)) + (1-t)g(\varphi(b)) - ct(1-t)(\varphi(a) - \varphi(b))^2.
\end{aligned}$$

Multiplying both sides of (2.10) by (2.11), it follows that

$$\begin{aligned}
& f(t\varphi(a) + (1-t)\varphi(b))g(t\varphi(a) + (1-t)\varphi(b)) \\
\leq & t^2 f(\varphi(a))g(\varphi(a)) + (1-t)^2 f(\varphi(b))g(\varphi(b)) \\
& + t(1-t)[f(\varphi(a))g(\varphi(b)) + f(\varphi(b))g(\varphi(a))] \\
& - ct^2(1-t)(\varphi(a) - \varphi(b))^2[f(\varphi(a)) + g(\varphi(a))] \\
& - ct(1-t)^2(\varphi(a) - \varphi(b))^2[f(\varphi(b)) + g(\varphi(b))] \\
& + c^2 t^2(1-t)^2(\varphi(a) - \varphi(b))^4.
\end{aligned}$$

By Lemma 1, it follows that the previous inequality can be integrated with respect to t over $[0, 1]$, getting

$$\begin{aligned}
& \int_0^1 f(t\varphi(a) + (1-t)\varphi(b))g(t\varphi(a) + (1-t)\varphi(b))dt \\
\leq & f(\varphi(a))g(\varphi(a)) \int_0^1 t^2 dt + f(\varphi(b))g(\varphi(b)) \int_0^1 (1-t)^2 dt \\
& + [f(\varphi(a))g(\varphi(b)) + f(\varphi(b))g(\varphi(a))] \int_0^1 t(1-t)dt \\
& - c(\varphi(a) - \varphi(b))^2[f(\varphi(a)) + g(\varphi(a))] \int_0^1 t^2(1-t)dt \\
& - c(\varphi(a) - \varphi(b))^2[f(\varphi(b)) + g(\varphi(b))] \int_0^1 t(1-t)^2 dt \\
& + c^2(\varphi(a) - \varphi(b))^4 \int_0^1 t^2(1-t)^2 dt.
\end{aligned}$$

In the first integral, we substitute $x = t\varphi(a) + (1-t)\varphi(b)$ and simple integrals calculated, we obtain the required inequality in (2.9). $\square$

REFERENCES

- [1] G. Cristescu and L. Lupşa, *Non-connected convexities and applications*, Kluwer Academic Publishers, Dordrecht / Boston / London, 2002.
- [2] G. Cristescu, *Hadamard type inequalities for φ -convex functions*, Annals of the University of Oradea, Fascicle of Management and Technological Engineering, CD-Rom Edition, III(XIII), 2004.
- [3] M. K. Bakula and J. Pečarić, *Note on some Hadamard-type inequalities*, Journal of Inequalities in Pure and Applied Mathematics, vol. 5, no. 3, article 74, 2004.

- [4] S. S. Dragomir and C. E. M. Pearce, *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University, 2000.
- [5] S. S. Dragomir and R.P. Agarwal, *Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula*, Appl. Math. lett., 11(5) (1998), 91-95.
- [6] J.E. Pečarić, F. Proschan and Y.L. Tong, *Convex Functions, Partial Orderings and Statistical Applications*, Academic Press, Boston, 1992.
- [7] E. Set, M. E. Özdemir, and S. S. Dragomir, *On the Hermite-Hadamard inequality and other integral inequalities involving two functions*, Journal of Inequalities and Applications, Article ID 148102, 9 pages, 2010.
- [8] E. Set, M. E. Özdemir, and S. S. Dragomir, *On Hadamard-Type inequalities involving several kinds of convexity*, Journal of Inequalities and Applications, Article ID 286845, 12 pages, 2010.
- [9] B.T. Polyak, *Existence theorems and convergence of minimizing sequences in extremum problems with restrictions*, Soviet Math. Dokl. 7 (1966), 72–75.
- [10] N. Merentes and K. Nikodem, *Remarks on strongly convex functions*, Aequationes Math. 80 (2010), no. 1-2, 193-199.
- [11] K. Nikodem and Zs. Pales, *Characterizations of inner product spaces by strongly convex functions*, Banach J. Math. Anal. 5 (2011), no. 1, 83–87.
- [12] H. Angulo, J. Gimenez, A. M. Moros and K. Nikodem, *On strongly h -convex functions*, Ann. Funct. Anal. 2 (2011), no. 2, 85–91.
- [13] E. A. Youness, *E - Convex Sets, E - Convex Functions and E - Convex Programming*, Journal of Optimization Theory and Applications, 102, 2(1999), 439-450.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, DÜZCE UNIVERSITY, DÜZCE-TURKEY

E-mail address: sarikayamz@gmail.com